

PERSPECTIVES IN MEASURING THE PPN PARAMETERS β and γ IN THE EARTH'S GRAVITATIONAL FIELD TO HIGH ACCURACY WITH CHAMP/GRACE MODELS

LORENZO IORIO

*Viale Unità di Italia 68, 70125 Bari (BA), Italy
e-mail: lorenzo.iorio@libero.it*

Received Day Month Year

Revised Day Month Year

Communicated by Managing Editor

The current bounds on the PPN parameters γ and β are of the order of $10^{-4} - 10^{-5}$. Various missions aimed at improving such limits by several orders of magnitude have more or less recently been proposed like LATOR, ASTROD, BepiColombo and GAIA. They involve the use of various spacecraft, to be launched along interplanetary trajectories, for measuring the post-Newtonian effects induced by the solar gravity on the propagation of electromagnetic waves. In this paper we investigate the requirements needed to measure the combination $\nu = (2 + 2\gamma - \beta)/3$ entering the post-Newtonian Einstein pericenter precession $\dot{\omega}$ of a test particle to an accuracy of the order of $\approx 10^{-5}$ with a pair of drag-free spacecraft in the Earth's gravitational field. It turns out that the latest gravity models from the dedicated CHAMP and GRACE missions would allow to reduce the systematic error of gravitational origin just to this demanding level of accuracy. In regard to the non-gravitational errors, the spectral noise density of the drag-free sensors required amounts to $10^{-8} - 10^{-9} \text{ cm s}^{-2} \text{ Hz}^{-\frac{1}{2}}$ over very low frequencies. Although not yet obtainable with the present technologies, such level of compensation is much less demanding than those required for, e.g., LISA. As a by-product, an independent measurement of the post-Newtonian gravitomagnetic Lense-Thirring effect with a $\approx 1\%$ accuracy would be possible as well. The forthcoming Earth gravity models from CHAMP and GRACE will further reduce the systematic bias of gravitational origin in both of such tests.

Keywords: PPN parameters γ and β ; drag-free spacecraft; CHAMP and GRACE Earth gravity models

1. Introduction

1.1. The PPN parameters γ and β

In the Parameterized Post-Newtonian (PPN) formalism¹, developed by Nordtvedt and Will to make easier the comparison of metric theories of gravity with each other and with experiment, the Eddington-Robertson-Schiff parameters β and γ describe how much non-linearity is present in the superposition law for gravity and how much spatial curvature is produced by a unit mass, respectively; in general relativity $\beta = \gamma = 1$. For a recent overview of the measurements of such PPN parameters see Refs. 2, 3.

The most accurate and clean determinations of γ come from the effects involving the propagation of electromagnetic signals: in their PPN expressions only γ is present, at first order. For the Shapiro time delay⁴, recently tested with the Doppler tracking of the Cassini spacecraft, the most accurate result is⁵ $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$, although the accuracy of such measurement has recently been somewhat questioned in Ref. 6; according to its authors, the motion of the Sun around the Solar System barycenter, if not properly accounted for, would induce a systematic bias of $\delta\gamma = 1.2 \times 10^{-4}$.

With β the situation is less favorable because it enters the post-Newtonian equations of motion of test particles in conjunction with γ itself in a rather complicated way. For example, in the post-Newtonian expression of the well known Einstein precession⁷ of the pericentre ω of a test particle the combination $\nu = (2 + 2\gamma - \beta)/3$ is present. E.V. Pitjeva in Ref. 8 recently processed more than 317 000 Solar System observations (1913-2003) for the construction of the Ephemerides of Planets and the Moon EPM2004 determining, among other things, β and γ with an accuracy of 10^{-4} . The same limit for β can also be obtained by combining the Cassini result for γ with the Lunar Laser Ranging test⁹ of the Nordtvedt^{10,11} effect^a, expressed in terms of $\eta = 4\beta - \gamma - 3$: indeed, from⁹ $\delta\eta = 7 \times 10^{-4}$, it follows $\delta\beta = 1.8 \times 10^{-4}$. Note that if $\delta\gamma = 1.2 \times 10^{-4}$ is assumed⁶, the error in β becomes $\delta\beta = 2.0 \times 10^{-4}$.

Various theoretical models involving scalar-tensor scenarios^{12,13,14,15,16,17} predicts deviations from unity for β and γ at a $10^{-6} - 10^{-7}$ level, so that it is very important to push the accuracy of such PPN tests towards this demanding accuracy.

The proposed LATOR mission¹⁸ has, among its goals, the determination of γ with an accuracy of 10^{-9} from the first order and a direct and independent measurement of β at a $\sim 0.01\%$ level via the second-order gravity-induced deflection of light. The expected accuracy in the proposed radio science experiments to be conducted with the future Mercury orbiter BepiColombo (<http://sci.esa.int/science-e/www/area/index.cfm?fareaid=30>) is of the order of¹⁹ 10^{-6} for β and γ . The accurate measurement of β and γ is one of the scientific goals also of the proposed mission ASTROD^{20,21}; the expected accuracy is²¹ 1×10^{-9} for both^b. The astrometric mission GAIA should be able to reach²³ $\delta\gamma \sim 10^{-7}$ through the deflection of light.

1.2. Aim of the paper

A satisfactorily empirical corroboration of a pillar of physics like general relativity requires that as many independent experiments as possible are conducted by different scientists in different laboratories. Now, all the performed or proposed high-precision tests of post-Newtonian gravity (in the weak-field and slow-motion approximation) are based on the effects induced by the gravitational field of the

^aIt is a violation of the strong equivalence principle which is zero in general relativity.

^bSee also Table 5 of Ref. 2 and Ref. 22.

Sun only. It is as if many independent experiments aimed to measure fundamental physical effects were conducted always in the same laboratory. Thus, it is worthwhile to try to use different laboratories, i.e. other gravitational fields, to perform such tests, even if their outcomes should be less accurate than those conducted in the Sun's field.

In this paper we will investigate if it is possible to realistically reach the $10^{-5} - 10^{-6}$ level of accuracy in measuring ν with a couple of dedicated drag-free spacecraft in the terrestrial gravitational field. This analysis is motivated by the fact that, in regard to the systematic errors of gravitational origin, reaching a 10^{-5} level would be right now possible thanks to the recent improvements in our knowledge of the classical part of the Earth's gravitational field due to the dedicated CHAMP (http://www.gfz-potsdam.de/pb1/op/champ/index_CHAMP.html) and GRACE (http://www.gfz-potsdam.de/pb1/op/grace/index_GRACE.html and <http://www.csr.utexas.edu/grace/>) missions. Much more demanding would be to reduce the non-gravitational errors down to the required level. As we will see, the drag-free technologies needed to implement such an ambitious goal are not yet available because they should work over observational time spans of several years; however, they would be less demanding than those required for complex missions like LISA²⁴.

2. The Einstein pericenter precession

The Einstein pericenter precession⁷ of a test particle in the gravitational field of a static central body is

$$\dot{\omega}_{\text{GE}} = \frac{3nGM}{c^2 a(1 - e^2)}, \quad (1)$$

where G is the Newtonian constant of gravitation, M is the mass of the central body which acts as source of the gravitational field, c is the speed of light in vacuum, a and e are the semimajor axis and the eccentricity, respectively, of the test particle's orbit and $n = \sqrt{GM/a^3}$ is its Keplerian mean motion.

As it is well known, Eq. (1) allowed to explain the Mercury's anomalous perihelion advance of 42.98 arcseconds per century observed since the 19th century. Modern radar-ranging measurements of the planetary perihelia in the solar field yielded a 10^{-3} accuracy^{25,26,27}.

Hiscock and Lindblom in Ref. 28 analyzed with some details the possibility of measuring the Einstein pericenter precessions of the natural satellites of Jupiter and Saturn which are much larger than those of the planets. Iorio in Ref. 29 showed that, at present, this possibility is not yet viable, despite of the first data sets from the Cassini mission.

The possibility of measuring the Einstein perigee rate in the Earth's gravity field with passive artificial satellites tracked via laser ranging was preliminarily investigated by Rubincam in Ref. 30 and Cugusi and Proverbio in Ref. 31. Ciufolini

and Matzner in Ref. 32 analyzed the data of the LAGEOS satellite using the GEM-L^{33,34} and GEM T1³⁵ Earth gravity models claiming a total accuracy of 20%, but such estimate is unrealistic both because of the too optimistic evaluation of the impact of the classical part of the Earth gravity field and of the treatment of the non-gravitational forces³⁶ acting on the perigee of the LAGEOS-like satellites. Moreover, the very small eccentricity of the LAGEOS orbit ($e = 0.0045$) makes its perigee badly defined and highly affected by the non-conservative perturbations. A more accurate and reliable test was proposed in Ref. 37 by suitably combining the nodes of LAGEOS and LAGEOS II and the perigee of LAGEOS II, and using the EGM96 Earth's gravity model³⁸. The estimated total accuracy is of the order of 10^{-3} . An analysis of the use of the perigee of LAGEOS II can be found in Ref. 39; the total error is estimated to be 2%. Also the proposed LAGEOS-like satellite LARES⁴⁰ could be used for measuring the Einstein perigee advance⁴¹. Hiscock and Lindblom in Ref. 28 envisaged the possibility of using geocentric drag-free spacecraft to measure the Einstein perigee precession. A step further concerning the use of dedicated terrestrial drag-free spacecraft for accurate tests of post-Newtonian gravity can be found in Ref. 42. In it the authors suggest that it would be possible to determine γ and β independently of each other with an accuracy of 10^{-4} by measuring ν and η .

3. Outline of the various sources of systematic errors

When a satellite-based test of a prediction of general relativity is proposed or performed, it is of the utmost importance to assess the total error budget in a very reliable and accurate way by accounting for all possible sources of systematic errors. Here we intend the aliasing action of other competing classical features of motion which may corrupt the genuine determination of the investigated relativistic effect. In regard to this topic, the choice of a suitable observable is very important in the sense that it must reach a good compromise between the errors of gravitational and non-gravitational origin whose simultaneous reduction is often in conflict.

3.1. The gravitational perturbations

The gravitational errors come from the mismodelling of the static and time-varying components of the multipolar expansion of the Newtonian gravitational potential of the Earth in spherical harmonics⁴³. Indeed, the static part of the even ($\ell = 2, 4, 6, \dots$) zonal ($m = 0$) harmonics $J_2, J_4, J_6, \dots$ induces secular precessions⁴³ on a satellite's node Ω and perigee ω

$$\begin{cases} \dot{\Omega}_{(\text{geopot})} = \sum_{\ell \geq 2} \dot{\Omega}_{\ell} J_{\ell}, \\ \dot{\omega}_{(\text{geopot})} = \sum_{\ell \geq 2} \dot{\omega}_{\ell} J_{\ell}, \end{cases} \quad (2)$$

where $\dot{\Omega}_{\ell}$ and $\dot{\omega}_{\ell}$ are the partial derivatives of the classical precessions with respect to the even zonal of degree ℓ (see, e.g., Ref. 44), while the odd zonal harmonics

$J_3, J_5, \dots$ affect the perigee with long-period harmonic perturbations having frequencies multiple of that of the perigee⁴³. The secular variations of the even zonal harmonics $\dot{J}_2, \dot{J}_4, \dot{J}_6, \dots$ induce quadratic signatures on the node and the perigee^{45,46}. There are also the long-period harmonic perturbations of the node and perigee due to the tides^{47,48,49} whose periods are determined by the lunisolar and satellite frequencies. The most insidious aliasing signals are the even zonal secular and quadratic advances and those harmonic signals that resemble superimposed linear trends over observational time spans of some years, like the $\ell = 2, m = 0$ constituent of the 18.6-year tide⁴⁷: their mismodeled part can corrupt to a significant extent the recovery of the genuine relativistic linear trends of interest^{48,49}.

A way to overcome the problem of the biasing classical even zonal secular trends consists in linearly combining the residuals $\delta\Psi$ of the Keplerian orbital elements, denoted generically as Ψ , affected by the classical and relativistic linear rates

$$\delta\dot{\Psi}^{(i)} = \sum_{\ell \geq 2} \dot{\Psi}_{\ell}^{(i)} \delta J_{\ell} + \dot{\Psi}_{\text{rel}}^{(i)} \xi_{\text{rel}}, \quad i = N, \quad (3)$$

where N is the number of perigees and/or nodes employed, in order to solve for the parameter ξ_{rel} , which accounts for the relativistic effect of interest ($\xi_{\text{rel}} = \nu$ in our case), and cancel out the impact of as many even zonal harmonics as possible among those recognized to be the most biasing ones^{50,51,52}. The coefficients $\dot{\Psi}_{\ell}^{(i)}$ are the partial derivatives of the classical precessions of $\Psi^{(i)}$ with respect to the even zonal of degree ℓ . The so-obtained linear combinations are sensitive to general relativity, irrespectively of the level of mismodelling δJ_{ℓ} of the static and time-varying parts of the selected even zonal harmonics. It turns out that the accuracy reached in the latest Earth gravity models from CHAMP and GRACE would allow right now to reduce the systematic error due to the geopotential to 10^{-5} of the Einstein precession if a suitable orbital geometry is chosen for a dedicated spacecraft-based mission. We discuss this topic in detail in Section 4.

3.2. The non-gravitational perturbations

On the other hand, such really important benefits cannot be fully capitalized if the non-gravitational forces^c acting on the spacecraft like the direct solar radiation pressure, Earth's albedo, Earth's IR radiation, atmospheric drag, thermal effects, etc.³⁶ are not suitably accounted for and reduced to the same level of the gravitational bias. For example, in the case of the Lense-Thirring⁵³ tests with the existing LAGEOS and the proposed LARES/OPTIS satellites^{40,54,55}, the obtainable accuracy is set up to $\sim 1\%$ by the non-conservative forces acting on the passive LAGEOS and LAGEOS II satellites which are not endowed with any active mechanism of compensation of such kind of perturbations^{56,57,58,59,60}; the systematic error due to the

^cLet us recall that their magnitude is proportional to the area-to-mass ratio σ/m of the satellite and depends on the interaction of the electromagnetic radiation with the satellite's physical structure, on its rotational state and its orbital geometry often in a very complex and intricate way.

geopotential would be much smaller according to the latest CHAMP/GRACE-based models⁶¹. Thus, it is apparent that only drag-free spacecraft could push the bias due to the non-conservative forces to the same level of that due to the gravitational perturbations. This topic will be the subject of Section 6.

4. How to reduce the gravitational perturbations

The idea of using a pair of drag-free Earth's artificial satellites was considered also in Refs. 62, 63, 64, 61; in that case the investigated feature of motion was the Lense-Thirring force which affects the node and the perigee of a satellite (see Section 8). In Refs. 62, 63, 64 it was shown that the difference of the perigees of two satellites in supplementary orbital configuration^d is able to add the Lense-Thirring precessions and to cancel out the classical even zonal precessions because of their functional dependence on the inclination⁴⁴. The use of active mechanisms of compensation of the non-gravitational perturbations was suggested because of the strong impact that they have on the perigees of LAGEOS-like satellites.

In the case of the Einstein precessions the situation is different since they are independent of the inclination. This implies that the difference of the perigees in the supplementary orbital configuration is not a good observable because it would exactly cancel the relativistic rates as well, independently of the values of γ and β . The linear combination approach involving the nodes and the perigees of both the spacecraft must, thus, be followed. Having the residuals of four measurable Keplerian orbital elements at our disposal it is possible to design a suitable observable

$$\delta\dot{\omega}^{(1)} + c_1\delta\dot{\omega}^{(2)} + c_2\delta\dot{\Omega}^{(1)} + c_3\delta\dot{\Omega}^{(2)} = \nu \left[\dot{\omega}_{\text{GE}}^{(1)} + c_1\dot{\omega}_{\text{GE}}^{(2)} \right] \quad (4)$$

which is affected by the Einstein precessions and is independent of the static and time-dependent parts of the first two even zonal harmonics of geopotential J_2, J_4 , and of the Lense-Thirring precessions. It turns out that, also in this case, the supplementary orbital configuration is not suitable because the coefficients weighing the perigees would be equal and opposite cancelling the relativistic signature. If different orbital parameters are chosen for the two spacecraft the situation becomes favorable. Indeed, with the orbital parameters of Table 1 we have for our

Table 1. Orbital parameters of the proposed drag-free spacecraft (1) and (2): a is in km and i in deg. Also their relativistic precessions, in mas yr⁻¹, are listed.

Satellite	a	i	e	$\dot{\omega}_{\text{GE}}$	$\dot{\omega}_{\text{LT}}$	$\dot{\Omega}_{\text{LT}}$
(1)	13000	50	0.2	2955.7	-53.0	27.5
(2)	12000	103	0.2	3610.5	23.6	35.0

^dIt means $a_1 = a_2$, $e_1 = e_2$, $i_1 = i_2 - 180$ deg. See Refs. 65, 66.

$J_2 - J_4$ –Lense-Thirring-free combination

$$\begin{cases} c_1 = 2.2817, \\ c_2 = -0.6861, \\ c_3 = 0.5165, \\ \dot{\omega}_{\text{GE}}^{(1)} + c_1 \dot{\omega}_{\text{GE}}^{(2)} = 11194.1 \text{ mas yr}^{-1}, \end{cases} \quad (5)$$

and the relative systematic error due to the uncanceled even zonal harmonics amounts to

$$\left. \frac{\delta\nu}{\nu} \right|_{\text{geopot}} \sim 3 \times 10^{-5}, \quad (6)$$

if the currently available CHAMP/GRACE-based EIGEN-CG03C Earth gravity model⁶⁷ is adopted for the calculation. Note that Eq. (6) represents a realistic upper bound of the gravitational bias because it has been obtained by linearly summing the absolute values of the individual errors up to degree $\ell = 20$. Note that this level of truncation is quite adequate because of the high altitudes of the satellites: indeed, it turns out that the numerical value of $\delta\nu/\nu|_{\text{geopot}}$ does not change if one adds more even zonals. The current models from CHAMP and GRACE are still preliminary and do not yet exploit the full data sets gathered by such missions which are still ongoing. Thus, it is likely that when new, more robust and reliable Earth gravity solutions will be available the estimate of Eq. (6) will further improve, all the satellites’ orbital parameters being equal.

In regard to the time-varying gravitational errors, the periods of the nodes and the perigees of the satellites are listed in Table 2. It can be noted that they amount

Table 2. Periods, in yr, of the node and the perigee of the proposed drag-free spacecraft (1) and (2)

Satellite	$P(\Omega)$	$P(\omega)$
(1)	-1.71	2.06
(2)	3.7	-2.22

to just a few years. This means that the sinusoidal perturbations induced on the nodes and the perigees by the $\ell = 2, m = 1$ constituent of the K_1 solar tide⁴⁷ and the odd zonal harmonic J_3 , which have just the same periods of the satellite’s node and perigee, respectively, and are not canceled out by the combination of Eq. (4), do not represent serious bias over reasonably long observational time spans. A potentially less favorable situation occurs for the $\ell = 3, m = 1, p = 1, q = -1$

oceanic constituent of the K_1 tide⁴⁷ which is not canceled by Eq. (4) and affects the perigees with perturbations whose periods are -10.0 yr and -5.6 yr, respectively. For the complete details of the action of the $\ell = 3$, $m = 1$ oceanic constituents of K_1 on the perigees of (1) and (2), see Table 3. According to the global ocean tide

Table 3. Periods P , in yr, and mismodeled amplitudes δA , in mas, of the tidal perturbations induced on the perigees of the proposed drag-free spacecraft (1) and (2) by the $\ell = 3$, $m = 1$ oceanic constituents of the K_1 solar tide according to the results of Table 6.4.8.5-1 of Ref. 36.

Satellite	$P(p = 1 \ q = -1)$	$\delta A(p = 1 \ q = -1)$	$P(p = 2 \ q = 1)$	$\delta A(p = 2 \ q = 1)$
(1)	-10.0	-10.1	-0.9	2.1
(2)	-5.6	7.0	1.4	3.3

model of the EGM96 solution³⁸, it turns out that the most insidious tidal effect is the K_1 $\ell = 3$, $m = 1$, $p = 1$, $q = -1$ perturbation on the perigee of satellite (1) with a mismodeled amplitude of -10.1 mas. Thus, the maximum relative error, over 10 yr, on the combination of Eq. (4) amounts to 9×10^{-5} . However, it must be noted that such an estimate is largely pessimistic both because it assumes a peak value after 10 yr and because it is based on the relatively old EGM96 model: more recent and accurate models like GOT99, CSR 4.0, FES2004 would greatly reduce the impact of such a source of bias.

If in constructing the combination of Eq. (4) one chose to cancel J_6 instead of the Lense-Thirring precessions, and a nearly-polar orbital configuration was adopted for both the satellites, a bound $\delta\nu/\nu|_{\text{geopot}} = 10^{-7}$ could be reached, according to EIGEN-CG03C. But, in this case, it would not be possible to separate^e the Einstein precessions from the Lense-Thirring ones which are weighted by a different combination of γ and β , namely $\mu = (1 + \gamma)/2$. Moreover, inclinations close to 90 deg would be harmful⁷⁰ because of the extremely long node periods and, thus, of the K_1 tidal perturbations.

5. The impact of the inclination errors

Another source of systematic error is related to the uncertainty in the knowledge of the spacecraft inclinations and their impact on the bias due to the uncanceled even zonal harmonics: indeed, recall that the coefficients $\dot{\omega}_\ell$ and $\dot{\Omega}_\ell$ of the classical even zonal precessions⁴⁴ do depend on the inclinations as well.

To this aim, we considered $\delta\nu/\nu|_{\text{geopot}}$ as a function of the two variables $i^{(1)}$ and $i^{(2)}$ and studied its behavior in small ranges $\Delta i^{(1)}$ and $\Delta i^{(2)}$ around the nominal values of Table 1 which simulate the errors in the inclinations. It turns out that for

^eAlso the post-Newtonian J_2/c^2 effects due to the Earth's oblateness^{68,69} affect all the combination proposed here. They must be modeled, otherwise they would induce a systematic bias of 10^{-4} on the measurement of the Einstein precession.

a very conservative and pessimistic assumption $\Delta i = 2$ deg for both $i^{(1)}$ and $i^{(2)}$ the relative error in ν varies by just 9×10^{-6} .

6. The accuracy required for the non-gravitational perturbations

In this Section we will investigate the level to which the non-gravitational accelerations should at most affect the spacecraft if a relative systematic error of 10^{-6} was to be obtained.

The Gauss equations for the variations of the node and the perigee are³⁶

$$\frac{d\Omega}{dt} = \frac{1}{na \sin i \sqrt{1-e^2}} A_n \left(\frac{r}{a} \right) \sin(\omega + f), \quad (7)$$

$$\frac{d\omega}{dt} = -\cos i \frac{d\Omega}{dt} + \frac{\sqrt{1-e^2}}{nae} \left[-A_r \cos f + A_t \left(1 + \frac{r}{p} \right) \sin f \right], \quad (8)$$

in which f is the true anomaly counted from the pericentre, $p = a(1-e^2)$ is the semilatus rectum of the Keplerian ellipse, i is the inclination of the orbit to the Earth's equator, A_r , A_t , A_n are the radial, transverse (in-plane components) and the normal (out-of-plane component) projections of the perturbing acceleration $\mathbf{A}$, respectively, on the co-moving frame $\{\hat{\mathbf{r}}, \hat{\mathbf{t}}, \hat{\mathbf{n}}\}$.

We are interested in the effects averaged over one orbital period P : they can be obtained by evaluating the right-hand-sides of Eq. (7) and Eq. (8) on the unperturbed Keplerian ellipse

$$r = \frac{p}{1 + e \cos f}, \quad (9)$$

and averaging them with

$$\frac{dt}{P} = \frac{(1-e^2)^{3/2}}{2\pi(1+e \cos f)^2} df. \quad (10)$$

Let us start with the node and consider the action of an out-of-plane acceleration consisting of a part $N^{(0)}$ constant over one orbital revolution and a part varying with the orbital frequency

$$A_n = N^{(0)} + N_S \sin f + N_C \cos f, \quad (11)$$

where N_S and N_C are to be considered as constant over one orbital revolution. Now Eq. (11) must be inserted into Eq. (7) and treated as previously outlined. By using

$$\begin{cases} \int_0^{2\pi} \frac{\sin(\omega+f)}{(1+e \cos f)^3} df = -\frac{3\pi e}{(1-e^2)^{5/2}} \sin \omega, \\ \int_0^{2\pi} \frac{\sin(\omega+f)(N_S \sin f + N_C \cos f)}{(1+e \cos f)^3} df = \frac{\pi}{(1-e^2)^{5/2}} [(1-e^2)N_S \cos \omega + (1+2e^2)N_C \sin \omega], \end{cases} \quad (12)$$

one finally gets

$$\left\langle \frac{d\Omega}{dt} \right\rangle = \frac{1}{2 \sin i} \sqrt{\frac{a}{GM(1-e^2)}} \{ [(1+2e^2)N_C - 3eN^{(0)}] \sin \omega + (1-e^2)N_S \cos \omega \}. \quad (13)$$

The same procedure holds for the perigee. By inserting

$$\begin{cases} A_r = R^{(0)} + R_S \sin f + R_C \cos f, \\ A_t = T^{(0)} + T_S \sin f + T_C \cos f \end{cases} \quad (14)$$

into the second term of the right-hand-side of Eq. (8) and using

$$\begin{cases} \int_0^{2\pi} \frac{\cos f}{(1+e \cos f)^2} df = -\frac{2\pi e}{(1-e^2)^{3/2}}, \\ \int_0^{2\pi} \frac{(2+e \cos f) \sin f}{(1+e \cos f)^3} df = 0, \\ \int_0^{2\pi} \frac{\cos f (R_S \sin f + R_C \cos f)}{(1+e \cos f)^2} df = \frac{2\pi R_C}{e^2} \left[1 - \frac{1-2e^2}{(1-e^2)^{3/2}} \right], \\ \int_0^{2\pi} \frac{(2+e \cos f)(T_S \sin f + T_C \cos f) \sin f}{(1+e \cos f)^3} df = -\frac{\pi T_S}{e^2} \left[2 + \frac{e^2-2}{(1-e^2)^{3/2}} \right], \end{cases} \quad (15)$$

it is possible to obtain

$$\left\langle \frac{d\omega}{dt} \right\rangle = -\cos i \left\langle \frac{d\Omega}{dt} \right\rangle + \left\langle \frac{d\omega}{dt} \right\rangle^{(0)} + \left\langle \frac{d\omega}{dt} \right\rangle^{(C/S)}, \quad (16)$$

with

$$\begin{cases} \left\langle \frac{d\omega}{dt} \right\rangle^{(0)} = \sqrt{\frac{a(1-e^2)}{GM}} R^{(0)}, \\ \left\langle \frac{d\omega}{dt} \right\rangle^{(C/S)} = -\sqrt{\frac{a}{GM}} \frac{(1-e^2)^2}{e^3} \left\{ \left[1 - \frac{1-2e^2}{(1-e^2)^{3/2}} \right] R_C + \left[1 + \frac{e^2-2}{2(1-e^2)^{3/2}} \right] T_S \right\}. \end{cases} \quad (17)$$

This result tells us that the perigee would be affected by secular or long-period perturbing rates, according to the frequencies embedded in $N^{(0)}$, N_S , N_T , $R^{(0)}$, R_C and T_S .

From Eq. (17) it is possible to obtain the level to which the non-gravitational perturbations should be reduced in order not to exceed a given systematic error in the measurement of the gravitoelectric perigee precession. For $a \sim 12000 - 13000$ km and $e = 0.2$, a 10^{-6} relative accuracy could be reached with a residual acceleration level of^f 10^{-11} cm s⁻². The drag-free technologies currently under development for, e.g., the LISA mission²⁴ should allow a cancelation of the non-gravitational accelerations down to 3×10^{-13} cm s⁻² Hz^{-1/2}. In our case, with an orbital frequency of 7×10^{-5} Hz, a drag-free level of $\sim 10^{-9}$ cm s⁻² Hz^{-1/2} would be sufficient: such a cancelation should be reached and maintained over time spans some years long, i.e. in a frequency range spanning from about 0 to 10^{-5} Hz. It is certainly not an easy task to reach. No long-term experience about the performance of such systems is available at present; current high precision accelerometers show-by extrapolating present data-a spectral noise density of⁷¹ $\sim 10^{-8}$ cm s⁻² Hz^{-1/2}. A very accurate in-orbit calibration would be required during the entire duration of the mission.

^fFor a comparison, the impact of the direct solar radiation pressure on the existing LAGEOS satellites amounts to⁵⁶ $\sim 10^{-7}$ cm s⁻².

7. The measurement errors

For the perigee and the node the measurable quantities are $r = ea\omega$ and $r = a \sin i\Omega$. Thus, for the combination of Eq. (4) the measurement error can be written as^g

$$\delta r_{\text{meas}} \left[\frac{1}{a^{(1)}e^{(1)}} + c_1 \frac{1}{a^{(2)}e^{(2)}} + c_2 \frac{1}{a^{(1)} \sin i^{(1)}} + c_3 \frac{1}{a^{(2)} \sin i^{(2)}} \right] \quad (18)$$

By using the values of Table 1 and by assuming $\delta r_{\text{meas}} = 0.1$ cm over an observational time span $T = 10$ yr, one gets for the relative error

$$\left. \frac{\delta \nu}{\nu} \right|_{\text{meas}} = 2 \times 10^{-6}. \quad (19)$$

About the assumptions made, it must be noted that we consider an experimental millimeter accuracy in determining the spacecraft orbit, in a root-mean-square sense, over 10 years: it is certainly not an easy task to be accomplished, especially in view of the fact that no empirical along-track accelerations should be fitted: otherwise the signal of interest could be removed as well. Moreover, it must be said that attempts to improve measurement errors are not the same as the capability of reconstructing the orbit with mm accuracy over 10 years.

8. The measurement of the Lense-Thirring effect

As a by-product, our active spacecraft could also be used to make an independent measurement of the Lense-Thirring precessions^{53,72,73,74,75}

$$\begin{cases} \dot{\Omega}_{\text{LT}} = \mu \frac{2GS}{c^2 a^3 (1-e^2)^{3/2}}, \\ \dot{\omega}_{\text{LT}} = -\mu \frac{6GS \cos i}{c^2 a^3 (1-e^2)^{3/2}}, \end{cases} \quad (20)$$

where S is the spin angular momentum of the central rotating mass. Indeed, if one combines the nodes and the perigees so to cancel out J_2 , J_4 and the Einstein precessions, and solves for μ , the resulting combination is obtained

$$\delta \dot{\omega}^{(1)} + k_1 \delta \dot{\omega}^{(2)} + k_2 \delta \dot{\Omega}^{(1)} + k_3 \delta \dot{\Omega}^{(2)} = \mu [\dot{\omega}_{\text{LT}}^{(1)} + k_1 \dot{\omega}_{\text{LT}}^{(2)} + k_2 \dot{\Omega}_{\text{LT}}^{(1)} + k_3 \dot{\Omega}_{\text{LT}}^{(2)}] \quad (21)$$

with

$$\begin{cases} k_1 = -0.9, \\ k_2 = 0.21, \\ k_3 = -2.7, \\ \dot{\omega}_{\text{LT}}^{(1)} + k_1 \dot{\omega}_{\text{LT}}^{(2)} + k_2 \dot{\Omega}_{\text{LT}}^{(1)} + k_3 \dot{\Omega}_{\text{LT}}^{(2)} = -161 \text{ mas yr}^{-1}. \end{cases} \quad (22)$$

^gPlease note that the labels (1) and (2) refer here to the two spacecraft.

In this case, the major limiting factor is represented by the systematic error due to the uncanceled even zonal harmonics because the drag-free level of cancelation of the non-gravitational perturbations discussed in Section 6 would allow to reduce their impact well below the percent level. It turns out that $\delta\mu/\mu|_{\text{geopot}} = 9 \times 10^{-3}$, according to EIGEN-CG03C. Such a result is not so good as it would be with the supplementary orbital configuration, and it could be reached also with only one new passive satellite and the existing LAGEOS and LAGEOS II (see Ref. 61). Nonetheless, $\delta\mu/\mu|_{\text{geopot}}$ should be improved by the new Earth gravity field solutions from CHAMP and GRACE pushing the overall accuracy of such test towards the $\sim 0.1\%$ level. This achievement could not be reached with the low-cost alternative involving one new satellite and the LAGEOS ones because of the uncanceled action of the non-gravitational perturbations.

9. Conclusions

In this paper we investigated what is needed to measure the combination $(2 + 2\gamma - \beta)/3$ of the post-Newtonian gravitoelectric Einstein perigee precession of a test particle at a $\approx 10^{-5}$ level of accuracy by using a couple of drag-free spacecraft in the Earth's gravitational field. The observable proposed is a linear combination of the nodes Ω and the perigees ω of such satellites to be analyzed over an observational time span of several years. It allows to cancel out the impact of the static and time-varying parts of the first two even zonal harmonics of the terrestrial geopotential, which represent a major source of systematic error, and of the post-Newtonian gravitomagnetic Lense-Thirring precessions. By using the latest Earth gravity model EIGEN-CG03C from CHAMP and GRACE, the bias induced by the remaining uncanceled even zonals amounts to 10^{-5} , for a suitable choice of the orbital configuration of the satellites with $a \sim 13000 - 12000$ km and $i \sim 50 - 103$ deg. When the forthcoming, more accurate and reliable CHAMP/GRACE-based Earth's gravity models will be available, it should be possible to push the gravitational bias down to 10^{-6} . An active reduction of the impact of the non-conservative perturbations is required in order to fully benefit of the low systematic error of gravitational origin. It turns out that the level of cancelation of the non-gravitational perturbations should be of the order of $10^{-9} \text{ cm s}^{-2} \text{ Hz}^{-\frac{1}{2}}$. Although the currently available drag-free sensors do not yet allow to reach such performances, also because they should be maintained over several years, they are less demanding than those required for, e.g., LISA. As a by-product of the proposed mission, an independent measurement of the Lense-Thirring effect with a $\approx 1\%$ accuracy would be possible as well. In this case, given the drag-free level required for the previous test, the most limiting factor is represented by the uncanceled even zonals; the forthcoming CHAMP/GRACE models should reduce their impact on this test as well. Another stringent and demanding requirement is reaching the ability of reconstructing the satellites' orbits at $\approx \text{mm}$ level in root-mean-square sense over a time span of 10 years.

References

1. C.M. Will, *Theory and Experiment in Gravitational Physics*, 2nd edition (Cambridge University Press, Cambridge, 1993).
2. W.-T. Ni, *Int. J. Mod. Phys. D* **14**, 901 (2005).
3. C.M. Will, *Living Rev. Relativity* **9**, 3 (2006). (www.livingreviews.org/lrr-2006-3)
4. I.I. Shapiro, *Phys. Rev. Lett.* **13**, 789 (1964).
5. B. Bertotti, L. Iess and P. Tortora, *Nature* **425**, 374 (2003).
6. S.M. Kopeikin *et al.*, *Phys. Lett. A* **367**, 276 (2007).
7. A. Einstein, *Sitzber. Kön. Preuss. Akad. Berlin* **47**, 831 (1915).
8. E.V. Pitjeva, *Astron. Lett.* **31**, 340 (2005).
9. J. Müller, J.G. Williams and S.G. Turyshev, Lunar Laser Ranging Contributions to Relativity and Geodesy, in *Lasers, Clocks, and Drag-Free: Technologies for Future Exploration in Space and Tests of Gravity*, eds. H. Dittus, C. Lämmerzahl and S.G. Turyshev (Springer-Verlag, Berlin, 2006), p. 357.
10. K. Nordtvedt, *Phys. Rev.* **169**, 1014 (1968a).
11. K. Nordtvedt, *Phys. Rev.* **169**, 1017 (1968b).
12. T. Damour and K. Nordtvedt, *Phys. Rev. Lett.* **70**, 2217 (1993a).
13. T. Damour and K. Nordtvedt, *Phys. Rev. D* **48**, 3436 (1993b).
14. T. Damour and G. Esposito-Farese, *Phys. Rev. D* **53**, 5541 (1996a).
15. T. Damour and G. Esposito-Farese, *Phys. Rev. D* **54**, 1474 (1996b).
16. T. Damour, F. Piazza and G. Veneziano, *Phys. Rev. Lett.* **89**, 081601 (2002a).
17. T. Damour, F. Piazza and G. Veneziano, *Phys. Rev. D* **66**, 046007 (2002b).
18. S.G. Turyshev, M. Shao and K. Nordtvedt, Science, Technology and Mission Design for the Laser Astrometric Test Of Relativity, in *Lasers, Clocks, and Drag-Free Control. Exploration of Relativistic Gravity*, eds. H. Dittus, C. Lämmerzahl and S.G. Turyshev (Springer-Verlag, Berlin, 2008), p. 473.
19. A. Milani *et al.*, *Phys. Rev. D* **66**, 082001 (2002).
20. W.-T. Ni, *Int. J. Mod. Phys. D* **11**, 947 (2002).
21. W.-T. Ni, S. Shiomi and A.-C. Liao, *Class. Quantum Grav.* **21**, S641 (2004).
22. W.-T. Ni, A. Pulido Patón and Y. Xia, Testing Relativistic Gravity to One Part per Billion, in *Lasers, Clocks, and Drag-Free Control. Exploration of Relativistic Gravity*, eds. H. Dittus, C. Lämmerzahl and S.G. Turyshev (Springer-Verlag, Berlin, 2008), p. 571.
23. A. Vecchiato *et al.*, *Astron. Astrophys.* **399**, 337 (2003).
24. D. Bortoluzzi *et al.*, *Class. Quantum Grav.* **20**, S89 (2003).
25. I.I. Shapiro *et al.*, *Phys. Rev. Lett.* **28**, 1594 (1972).
26. I.I. Shapiro, C. Counselman and R. King, *Phys. Rev. Lett.* **36**, 555 (1976).
27. I.I. Shapiro, Solar system tests of general relativity: recent results and present plans, in *Proceedings of the 12th International Conference on General Relativity and Gravitation (Boulder, Colorado, 1989)*, eds. N. Ashby, D. Bartlett and W. Wyss (Cambridge University Press, Cambridge, 1990), p. 313.
28. W.A. Hiscock and L. Lindblom, *Astroph. J.* **231**, 224 (1979).
29. L. Iorio, *Int. J. Mod. Phys. D* **16**, 11 (2007).
30. D.P. Rubincam, *Celest. Mech. Dyn. Astron.* **15**, 21 (1977).
31. L. Cugusi and E. Proverbio, *Astron. Astrophys.* **69**, 321 (1978).
32. I. Ciufolini and R. Matzner, *Int. J. Mod. Phys. A* **7**, 843 (1992).
33. F.J. Lerch *et al.*, *J. Geophys. Res.* **90**(B11), 9301 (1985a).
34. F.J. Lerch *et al.*, *J. Geophys. Res.* **90**(B11), 9312 (1985b).
35. J.G. Marsh *et al.*, *J. Geophys. Res.* **93**(B6), 6169 (1988).
36. A. Milani, P. Farinella and A.M. Nobili, *Non-Gravitational Perturbations and Satellite*

- Geodesy* (Adam Hilger, Bristol, 1987).
37. L. Iorio, I. Ciufolini and E.C. Pavlis, *Class. Quantum Grav.* **19**, 4301 (2002a).
 38. F.G. Lemoine *et al.*, NASA/TP-1998-206861, (1998).
 39. D.M. Lucchesi, *Phys. Lett. A* **318**, 234 (2003a).
 40. I. Ciufolini *et al.*, *LARES Phase A* (University La Sapienza, Rome, 1998).
 41. L. Iorio, D.M. Lucchesi and I. Ciufolini, *Class. Quantum Grav.* **19**, 4311 (2002b).
 42. T. Damour and G. Esposito-Farese, *Phys. Rev. D* **50**, 2381 (1994).
 43. W.M. Kaula, *Theory of Satellite Geodesy* (Dover, 2000).
 44. L. Iorio, *Celest. Mech. Dyn. Astron.* **86**, 277 (2003a).
 45. D.M. Lucchesi, *Int. J. Mod. Phys. D* **14**, 1989 (2005).
 46. L. Iorio, *J. of Geodesy* **80**, 128 (2006).
 47. L. Iorio, *Celest. Mech. Dyn. Astron.* **79**, 201 (2001a).
 48. L. Iorio and E.C. Pavlis, *J. of the Geod. Soc. of Japan* **47**, 169 (2001).
 49. E.C. Pavlis and L. Iorio, *Int. J. Mod. Phys. D* **11**, 599 (2002).
 50. I. Ciufolini, *Nuovo Cim. B* **109**, 1709 (1996).
 51. L. Iorio, *Class. Quantum Grav.* **19**, 5473 (2002a).
 52. L. Iorio, Possible New Approaches for Improving the Accuracy in the Lense-Thirring Effect Measurement, in *Horizons in World Physics*, Vol. 245, ed. T.V. Lynch (Nova, Hauppauge, 2004), p. 273.
 53. J. Lense and H. Thirring, *Phys. Z.* **19**, 156 (1918). Translated and discussed by B. Mashhoon, F.W. Hehl and D.S. Theiss, *Gen. Relativ. Grav.* **16**, 711 (1984). Reprinted in R.J. Ruffini and C. Sigismondi (eds.), *Nonlinear Gravitodynamics* (World Scientific, Singapore, 2003), p. 349.
 54. L. Iorio *et al.*, *Class. Quantum Grav.* **21**, 2139 (2004).
 55. C. Lämmerzahl *et al.*, *Gen. Relativ. Grav.* **36**, 2373 (2004).
 56. D.M. Lucchesi, *Plan. Space Sci.* **49**, 447 (2001).
 57. D.M. Lucchesi, *Plan. Space Sci.* **50**, 1067 (2002).
 58. D.M. Lucchesi, *Geophys. Res. Lett.* **30**, 1957 (2003b).
 59. D.M. Lucchesi, *Celest. Mech. Dyn. Astron.* **88**, 269 (2004).
 60. D.M. Lucchesi *et al.*, *Plan. Space Sci.* **52**, 699 (2004).
 61. L. Iorio, *New Astron.* **10**, 616 (2005).
 62. L. Iorio, *Phys. Lett. A* **308**, 81 (2003b).
 63. L. Iorio, *Gen. Rel. Gravit.* **35**, 1583 (2003c).
 64. L. Iorio and D.M. Lucchesi, *Class. Quantum Grav.* **20**, 2477 (2003).
 65. I. Ciufolini, *Phys. Rev. Lett.* **56**, 278 (1986).
 66. J.C. Ries, R.J. Eanes, M.M. Watkins and B.D. Tapley, *Joint NASA/ASI Study on Measuring the Lense-Thirring Precession Using a Second LAGEOS Satellite*, CSR-89-3 (Center for Space Research, The University of Texas at Austin, 1989).
 67. C. Förste, *et al.*, Poster g004_EGU05-A-04561.pdf presented at *EGU General Assembly 2005 (Vienna, Austria, 24-29 April)* (2005).
 68. M.H. Soffel *et al.*, *Celest. Mech. Dyn. Astron.* **42**, 81 (1988).
 69. J. Heimberger, M.H. Soffel and H. Ruder, *Celest. Mech. Dyn. Astron.* **47**, 205 (1990).
 70. L. Iorio, *Class. Quantum Grav.* **19**, L175 (2002b).
 71. S. Scheithauer *et al.*, *Aerospace Science and Technology* **9**, 357 (2005).
 72. B.M. Barker and R.F. O'Connell, *Phys. Rev. D* **10**, 1340 (1974).
 73. M.H. Soffel, *Relativity in Astrometry, Celestial Mechanics and Geodesy* (Springer-Verlag, Berlin, 1989).
 74. N. Ashby and T. Allison, *Celest. Mech. Dyn. Astron.* **57**, 537 (1993).
 75. L. Iorio, *Nuovo Cim. B* **116**, 777 (2001b).